



**QUEEN'S  
UNIVERSITY  
BELFAST**

## **Statin Use and Survival from Lung Cancer: A Population-Based Cohort Study**

Cardwell, C. R., McMenamin, Ú., Hughes, C. M., & Murray, L. J. (2015). Statin Use and Survival from Lung Cancer: A Population-Based Cohort Study. *Cancer Epidemiology Biomarkers & Prevention*, 24(5), 833-841. <https://doi.org/10.1158/1055-9965.EPI-15-0052>

**Published in:**  
Cancer Epidemiology Biomarkers & Prevention

**Document Version:**  
Peer reviewed version

**Queen's University Belfast - Research Portal:**  
[Link to publication record in Queen's University Belfast Research Portal](#)

**Publisher rights**  
Copyright 2015 The American Association for Cancer Research (AACR), Inc.

**General rights**  
Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**  
The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact [openaccess@qub.ac.uk](mailto:openaccess@qub.ac.uk).

**Title:** Statin use and survival from lung cancer: a population-based cohort study.

**Authors and affiliations:** Chris R Cardwell<sup>1</sup>, Úna Mc Menamin<sup>1</sup>, Carmel M Hughes<sup>2</sup>, Liam J Murray<sup>1,3</sup>.

<sup>1</sup>Cancer Epidemiology and Health Services Research Group, Centre for Public Health, Queen's University Belfast, Belfast, Northern Ireland.

<sup>2</sup>School of Pharmacy, Queen's University Belfast, Northern Ireland, BT9 7BL.

<sup>3</sup>Centre of Excellence for Public Health (NI), Centre for Public Health, Queen's University Belfast, Belfast, Northern Ireland.

**Running title:** Statin use and survival from lung cancer.

**Key words:** Lung cancer, statins, mortality, adjunct treatment, epidemiology.

**Financial support:** CRC and UM were supported by a Health and Social Care Research and Development, Public Health Agency, Northern Ireland, funded UK NIHR Career Development Fellowship, which also funded access to the CPRD dataset. The funders had no role in the study design; collection, analysis, and interpretation of data; writing of the report; or the decision to submit for publication.

**Corresponding author:** Dr Chris Cardwell ([c.cardwell@qub.ac.uk](mailto:c.cardwell@qub.ac.uk)), Institute of Clinical Sciences Block B, Queen's University Belfast, Royal Victoria Hospital, Grosvenor Road, Belfast, BT12 6BA. Phone: +44(0)28 90632620. Fax: +44(0)28 90235900.

**Conflicts of interest:** The authors have no conflicts of interest to disclose.

**Word count:** 2,726

**Number of figures:** 0.

**Number of Tables:** 5.

## ABSTRACT

**Background:** Preclinical evidence from lung cancer cell lines and animal models suggest that statins could have anticancer properties. We investigated whether statin users had reduced risk of cancer-specific mortality in a population based cohort of lung cancer patients.

**Methods:** Newly diagnosed lung cancer patients, from 1998 to 2009, were identified from English cancer registry data and linked to the UK Clinical Practice Research Datalink, providing prescription records, and to Office of National Statistics mortality data up to 2012. Cox regression models were used to calculate hazard ratios (HR) for cancer-specific mortality and 95% confidence intervals (CIs) by statin use before and after diagnosis and to adjust these HRs for potential confounders.

**Results:** In 3,638 lung cancer patients, there was some evidence that statin use after diagnosis was associated with reduced lung cancer-specific mortality (adjusted HR=0.89, 95% CI 0.78, 1.02; P=0.09). Associations were more marked after 12 prescriptions (adjusted HR=0.81, 95% CI 0.67, 0.98; P=0.03) and when lipophilic statins were investigated (adjusted HR=0.81, 95% CI 0.70, 0.94; P=0.01) but were attenuated in some sensitivity analyses. Furthermore, in 11,051 lung cancer patients, statin use before diagnosis was associated with reduced lung cancer-specific mortality (adjusted HR=0.88, 95% CI, 0.83, 0.93; P<0.001).

**Conclusions:** There was some evidence that lung cancer patients who used statins, and particularly simvastatin, had reduced rates of cancer-specific mortality.

**Impact:** These findings should first be confirmed in observational studies, but provide some support for conducting randomized controlled trials of simvastatin as adjuvant cancer therapy in lung cancer patients.

## Introduction

Statins (3-hydroxy-3-methylglutaryl coenzyme A reductase inhibitors) are primarily used for primary and secondary prevention of coronary heart disease.(1) In recent decades the potential anticancer properties of statins have received much attention.(2-4) Preclinical studies have proposed a number of relevant biological mechanisms including inhibition of cell proliferation, stimulation of anticancer immune surveillance, and interruption of oncogenic signalling.(2) In humans, epidemiological studies have shown reductions in cancer recurrence or cancer-specific mortality at a number of cancer sites including breast(5), prostate(6) and colorectal.(7)

The preclinical evidence for the effect of statins on lung cancer appears promising. In vitro studies have shown reduced proliferation(8, 9), reduced migration(10) and increased apoptosis(9) in lung cancer cell lines treated with simvastatin and an in vivo study,(10) using a mouse model, demonstrated reduced tumour growth and bone metastases after simvastatin treatment. Despite this preclinical evidence, few studies in humans have investigated the association between statin use and survival in lung cancer patients. A recent Danish study(11) observed some evidence of a weak reduction in cancer-specific mortality in individuals who used statins before lung cancer diagnosis (HR=0.87 95% CI 0.83, 0.92). However, this study did not report findings for statins used after diagnosis, the time period when clinical intervention is possible, nor did it report results for lung cancer by statin type or evidence of dose response associations. A phase 3 randomised clinical trial has also been conducted in 846 small cell lung cancer patients which compared chemotherapy and pravastatin to chemotherapy alone.(12) Although the final trial report has not yet been published, the results were presented at the World Conference on Lung Cancer, Sydney, and a significant protective effect was not observed.(13) It is worth noting that this trial only investigated pravastatin and included only patients with small cell lung cancer. Consequently, there

remains a need to investigate the association between statin use and survival after lung cancer diagnosis, particularly for simvastatin because of the considerable supportive preclinical evidence.

Therefore, we investigated the effect of statin use, both before and after cancer diagnosis, on cancer-specific mortality in a large population-based cohort of lung cancer patients.

## **Materials and Methods**

### **Data source**

This study is based upon linkages between the National Cancer Data Repository (NCDR), the UK Clinical Practice Research Datalink (CPRD) and the Office of National Statistics (ONS) death registration data. NCDR data, available for England, included all patients identified by cancer registries, containing date and site of primary cancer diagnosis, and treatment data.

The CPRD includes general practice records and contains demographic information, clinical diagnoses, and prescription data of documented high quality(14-16). CPRD is the world's largest computerized database of longitudinal patient records comprising approximately 7% of the UK population. ONS death registration data provided date and cause of death.

Linkages between data sources were conducted using a deterministic algorithm based upon NHS number, gender, date of birth, and postcode. Ethical approval for all observational research using CPRD data has been obtained from a multicentre research ethics committee.

### **Cohort selection**

A cohort of newly diagnosed lung cancer patients was identified from cancer registry recorded primary diagnosis of lung cancer (ICD code C34 Malignant neoplasm of bronchus and lung) between 1998 and 2009, from NCDR. Cohort members with previous NCDR cancer diagnosis, apart from in situ neoplasms and non-melanoma skin cancers, were excluded. This cohort was linked to CPRD and ONS death registration data. Deaths were identified from ONS with coverage up to January 2012 with lung cancer-specific deaths defined as those with underlying cause of death ICD code C34.

### **Exposure data**

Statins were determined from GP prescribing records (from CPRD) based upon the Statins section of the British National Formulary (Section 2.12).(17) A quantity of 28 tablets (the

most common pack size prescribed) was assumed for approximately 1% of prescriptions where quantity was missing or assumed incorrect. The daily defined doses (DDD) in each prescription were calculated by multiplying the quantity by the strength (in mg) and dividing by the mg in a DDD from the World Health Organisation.(18) In the main analysis of statins after diagnosis, the exposure window for statin use was from lung cancer diagnosis to death or end of follow-up, but was subject to a lag (described later). In the main analysis of statins before diagnosis the exposure window was from one year prior to lung cancer diagnosis to the date of lung cancer diagnosis. Statins were analysed individually and grouped by type into lipophilic statins (including simvastatin, fluvastatin and cerivastatin) and hydrophilic statins (including atorvastatin, pravastatin and rosuvastatin).

### **Covariates**

Data available from NCDR included histology, surgery, chemotherapy and radiotherapy in the six months after diagnosis. Cancer registry recorded histology codes (International Classification of Diseases for Oncology, 3<sup>rd</sup> edition) were used to identify small cell lung cancer patients. Smoking, alcohol, and body mass index (BMI) were determined from the closest GP record prior to lung cancer diagnosis (values more than ten years prior to diagnosis were ignored). Comorbidities were determined prior to diagnosis from GP diagnosis codes using comorbidities comprising a recent adaptation of the Charlson index.(19) CPRD provided deprivation based upon postcode of residence using the 2004 index of multiple deprivation for England.(20) Low-dose aspirin, beta-blocker use and bisphosphonate use were determined from GP prescription records to allow for adjustment, because of potential associations with cancer-specific mortality.

### **Statistical analysis**

In analyses of statin use after lung cancer diagnosis, deaths in the first year after diagnosis were removed as it seemed unlikely that statin use after diagnosis could influence such deaths. Consequently, patients were followed from one year after lung cancer diagnosis to death, end of registration with the general practice, last date of data collection from general practice or end of ONS follow-up. Time-dependent Cox regression models were used to calculate hazard ratios for lung cancer-specific death (HRs) and 95% confidence intervals (95% CIs). Statin use was defined as a time varying covariate(21) with patients initially considered non-users and then users after a lag of 6 months after their first statin prescription. A lag is recommended(22) and was used to remove prescriptions in the 6 months prior to death as these may reflect increased contact with medical professionals or end of life care. In sensitivity analyses the duration of this lag was varied. Dose-response analyses were conducted with individuals considered non-users prior to 6 months after first use, a short term user between 6 months after first use and 6 months after the 12<sup>th</sup> prescription (or 365<sup>th</sup> DDD) and a longer term user after this time. Adjusted analyses were conducted by including potential confounders in the cox regression models. Analyses were conducted by number of prescriptions, number of DDDs and type of statin and repeated for all-cause mortality (which does not require classification of the cause of death). Subgroup analyses were conducted by gender, type of lung cancer (i.e. small cell versus non-small cell) and pre-diagnostic statin use. Tests for interactions were performed using interaction terms within the Cox regression models. Sensitivity analyses were conducted increasing the lag to 1 year, additionally adjusting for smoking, BMI, bisphosphonate use and lung cancer type. A simplified analysis was conducted comparing statin users to non-users in the first year after lung cancer diagnosis in individuals alive 1 year after diagnosis, thus controlling for immortal time bias(23) without requiring time varying covariates. A nested case-control analysis of the cohort was also conducted in which cases who died due to lung cancer were matched on gender, age (in 5 years) and year of diagnosis (in 2 years) to five risk-set controls who lived



at least as long after their lung cancer diagnosis. The exposure period was from lung cancer diagnosis until 6 months prior to lung cancer-specific death in cases and for a period of identical duration from diagnosis in matched controls. Conditional logistic regression was then used to calculate corresponding odds ratios (ORs), and 95% confidence intervals (CIs), for statin use in the exposure period adjusting for potential confounders.

In analyses of statin use before cancer diagnosis, lung cancer patients were followed from diagnosis to death or censoring (as defined previously). Deaths in the first year after diagnosis were not excluded. Cox regression models were used to calculate HRs and 95% CIs for statin use based upon prescriptions in the year prior to diagnosis (restricted to individuals with at least 1 year of records prior to diagnosis). Adjusted analyses were conducted including only potential confounders recorded prior to diagnosis to avoid over-adjustment (such as age, year, gender, comorbidities, deprivation, aspirin and beta-blocker user in the year prior to diagnosis).<sup>(24, 25)</sup> Analyses were conducted by statin type, number of prescriptions and number of DDDs. Subgroup analyses were conducted by gender. Sensitivity analyses were conducted additionally adjusting for BMI, bisphosphonate use and smoking prior to diagnosis and defining statin use as a prescription from 2 years to 6 months prior to diagnosis (restricted to individuals with at least 2 years of records prior to diagnosis).

## **Results**

### **Patient cohort**

A total of 14,689 lung cancer patients were diagnosed in NCDR with linked CPRD data. The analysis of statin use after lung cancer contained 3,638 patients, because 11,051 were excluded with less than 1 year of follow-up (10,265 of whom had died), with average follow-up of 3 years (maximum 14 years). The analysis of statin use before lung cancer contained 13,398 patients, because 1,291 were excluded with less than 1 year of records before diagnosis, with average follow-up of 1 year (maximum 14 years).

Patient characteristics by statin use are shown in Table 1. Statin users (before or after diagnosis) were more likely to be diagnosed more recently, be older, be male, have higher BMI, have comorbidities (particularly for cerebrovascular disease, diabetes and myocardial infarction) and to use aspirin and beta-blockers. Statin users after diagnosis were less likely to have chemotherapy. The associations between statin use and other patient characteristics were less marked. In 3,357 lung cancer patients who lived at least 1 year after diagnosis and who had 1 year of records before diagnosis, 2,223 did not use statins in the year before diagnosis and did not use statins any time after diagnosis, 76 used before but not after, 256 used after but not before and 802 used before and after.

### **Statin use after diagnosis**

Overall, statin users after diagnosis had an 11% reduction in the rate of lung cancer-specific mortality (adjusted HR=0.89, 95% CI 0.78, 1.02; P=0.09) compared with statin non-users after adjustment for confounders, Table 2. Compared with statin non-users, individuals with 1 to 12 statin prescriptions after diagnosis had a 6% reduction (adjusted HR=0.94, 95% CI 0.81, 1.09; P=0.39) whilst individuals with more than 12 prescriptions had a 19% reduction (adjusted HR=0.81, 95% CI 0.67, 0.98; P=0.03) in the rate of lung cancer-specific mortality.

A similar pattern was seen for statin use based upon DDDs. There was some indication that lipophilic statins (adjusted HR=0.81, 95% CI 0.70, 0.94; P=0.01), particularly simvastatin (adjusted HR=0.80, 95% CI 0.69, 0.93; P=0.003), appeared to have more protective associations than hydrophilic agents (adjusted HR=1.00, 95% CI 0.85, 1.18; P=0.01). Similar associations were observed for all-cause mortality. In subgroup analyses (Table 3) there were no marked differences in associations by gender (P for interaction=0.47), pre-diagnostic statin use (P for interaction=0.49) or by lung cancer type (P for interaction=0.12). The associations were little altered in most sensitivity analyses but they were attenuated when the lag was increased to 1 year (adjusted HR=0.92, 95% CI 0.80, 1.05; P=0.22)

### **Statin use before diagnosis**

Overall, statin users before diagnosis had a 12% reduction in rate of lung cancer-specific mortality (adjusted HR=0.88, 95% CI 0.83, 0.93; P<0.001) compared with statin non-users following adjustment for confounders, Table 4. The magnitude of the reduction in risk of lung cancer-specific death was similar in patients with 1 to 12 statin prescriptions in the year before diagnosis (adjusted HR=0.87, 95% CI 0.82, 0.92; P<0.001) and in those with more than 12 prescriptions (adjusted HR=0.91, 95% CI 0.84, 0.99; P=0.02) compared with statin non-users. Comparable associations were observed for all-cause mortality. The observed associations were similar by statin type, by gender (P for interaction=0.10) and in sensitivity analyses, Table 5.

## Discussion

In this large population-based cohort, lung cancer patients who used statins after diagnosis had a non-significant 11% reduction in cancer-specific mortality and those who used simvastatin had a 20% reduction in cancer-specific mortality. These associations were apparent when prescriptions in the 6 months prior to death, which could reflect end of life care, were removed but were attenuated in a sensitivity analysis which removed prescriptions in the year prior to death. Lung cancer patients who used statins in the year before diagnosis had a 12% reduction in cancer-specific mortality. This association was little altered in sensitivity analyses.

Although there have not been any epidemiological studies which have investigated the association between statin use after lung cancer diagnosis and cancer-specific mortality, a recent trial(12, 13) found no protective effect of pravastatin and chemotherapy compared to chemotherapy alone in small cell lung cancer patients. Our study also found no association between pravastatin use after diagnosis and cancer specific mortality (adjusted HR=1.23 95% CI 0.85, 1.77; P=0.27), but we found some evidence of protective associations for other statins which were not investigated in this trial, such as simvastatin. A similar phase 2 trial(26) has also been conducted which included 106 advanced (stage 3b or 4) non-small cell lung cancer patients. This trial observed a weak improvement in survival in a group that received simvastatin plus gefitinib compared with gefitinib only (HR=0.88 95% CI 0.57, 1.35; P=0.49), but this was not significant. We observed a slightly more marked, but not inconsistent, inverse association between simvastatin and cancer-specific mortality (adjusted HR=0.80 95% CI 0.69, 0.93; P=0.003). Other early phase trials, though not powered to report upon efficacy, are also in progress(27-30) but contain relatively small numbers.

Statin use before diagnosis and survival of lung cancer patients was investigated in a subgroup of a large Danish epidemiological study.(11) This Danish study demonstrated a weak protective effect of pre-diagnostic statins on cancer-specific mortality in lung cancer patients (adjusted HR=0.87 95% CI 0.83, 0.92) almost identical to our finding (adjusted HR=0.88 95% CI 0.83, 0.93), but this study did not report results for different types of statins or dose response analyses in lung cancer patients.

Our study has several strengths and limitations. This is the first study to investigate statin use after diagnosis and lung cancer-specific mortality. The cohort was large with long follow-up of up to 14 years. NCDR and ONS data allowed robust verification of cancer diagnosis and death. Although some misclassification of cause of death is possible, this seems unlikely to bias results particularly as analysis of all-cause mortality produced similar findings. Routinely recorded GP dates of prescriptions were used eliminating the potential for recall bias. However, misclassification of statin use is possible due to over the counter use, but only low dose 10mg simvastatin is available over the counter in the UK and only from 2004.(31) Non-compliance may also contribute to misclassification, but associations were also apparent in patients who had 12 or more statin prescriptions in whom non-compliance is of less concern. Furthermore, any misclassification of statin usage is likely to drag associations to the null rather than create spurious associations.

The cause of any reduction in lung cancer-specific mortality in lung cancer patients using statins is unknown. As with all observational studies, it is not possible to rule out residual confounding by unrecorded or incomplete variables. In particular adjustments for stage were not possible which could influence results for statins after diagnosis but such adjustments are probably not appropriate for the analysis of statins before diagnosis, as stage may lie on the causal pathway. The observed associations could also be influenced by confounding by

indication(32) or the healthy user effect,(33) though these biases would assume that statin users were healthier but in our cohort statin users had higher BMI and more comorbidities. The observed associations are consistent with the pre-clinical evidence for anti-cancer properties of statins from lung cancer cell lines(8-10) and mouse models.(10) These results require confirmation in further large epidemiological studies, particularly those with complete stage data, which could inform the decision to conduct a trial of simvastatin in lung cancer patients.

In conclusion, in this large population-based lung cancer cohort, there were weak inverse associations between statin usage and time to cancer-specific death. These associations could be causal or could reflect residual confounding or other biases.

## **Acknowledgements**

This study is based partly on data from the CPRD obtained under license from the UK Medicines and Healthcare products Regulatory Agency. However, the interpretation and conclusions contained in this study are those of the authors alone. None of the authors have any conflicts of interest to declare.

## References

- (1) Boudreau DM, Yu O, Johnson J. Statin use and cancer risk: a comprehensive review. *Expert Opin Drug Saf* 2010;9:603-621.
- (2) Ahern TP, Lash TL, Damkier P, Christiansen PM, Cronin-Fenton DP. Statins and breast cancer prognosis: evidence and opportunities. *Lancet Oncol* 2014;15:e461-e468.
- (3) Chan KK, Oza AM, Siu LL. The statins as anticancer agents. *Clin Cancer Res* 2003;9:10-9.
- (4) Vallianou NG, Kostantinou A, Kougias M, Kazazis C. Statins and cancer. *Anticancer Agents Med Chem* 2014;14:706-12.
- (5) Ahern TP, Pedersen L, Tarp M, Cronin-Fenton DP, Garne JP, Silliman RA et al. Statin prescriptions and breast cancer recurrence risk: a Danish nationwide prospective cohort study. *J Natl Cancer Inst* 2011;103:1461-1468.
- (6) Park HS, Schoenfeld JD, Mailhot RB, et al. Statins and prostate cancer recurrence following radical prostatectomy or radiotherapy: a systematic review and meta-analysis. *Ann Oncol* 2013;24:1427-34.
- (7) Cardwell CR, Hicks BM, Hughes C, Murray LJ. Statin use after colorectal cancer diagnosis and survival: a population-based cohort study. *J Clin Oncol* 2014;32:3177-3183.
- (8) Li Y, Fu J, Yuan X, Hu C. Simvastatin inhibits the proliferation of A549 lung cancer cells through oxidative stress and up-regulation of SOD2. *Pharmazie* 2014;69:610-614.
- (9) Yu X, Pan Y, Ma H, Li W. Simvastatin inhibits proliferation and induces apoptosis in human lung cancer cells. *Oncol Res* 2013;20:351-357.
- (10) Liu H, Wang Z, Li Y, Li W, Chen Y. Simvastatin prevents proliferation and bone metastases of lung adenocarcinoma in vitro and in vivo. *Neoplasma* 2013;60:240-246.
- (11) Nielsen SF, Nordestgaard BG, Bojesen SE. Statin use and reduced cancer-related mortality. *N Engl J Med* 2012;367:1792-1802.



- (12) Seckl M, Ottensmeier C, Cullen M, et al. A multicenter phase III randomized double-blind placebo controlled trial of pravastatin added to first-line standard chemotherapy in patients with small cell lung cancer (SCLC). J Clin Oncol 2013;31(Suppl; Abstract 7595).
- (13) Statins do not extend survival of SCLC patients on chemotherapy. 2013. 15th World Conference on Lung Cancer News Wednesday October 30, 2013 [Internet]. International Association for the Study of Lung Cancer [cited 2014 Nov 1]. Available from <http://www.2013worldlungcancer.org/press-media/documents/WCLC2013-Day4-Newspaper.pdf>.
- (14) Jick H, Jick SS, Derby LE. Validation of information recorded on general practitioner based computerised data resource in the United Kingdom. BMJ 1991;302:766-768.
- (15) Boggon R, van Staa TP, Chapman M, et al. Cancer recording and mortality in the General Practice Research Database and linked cancer registries. Pharmacoepidemiol Drug Saf 2013;22:168-75.
- (16) Herrett E, Thomas SL, Schoonen WM, Smeeth L, Hall AJ. Validation and validity of diagnoses in the General Practice Research Database: a systematic review. Br J Clin Pharmacol 2010;69:4-14.
- (17) British National Formulary 66th ed. 2013. Joint Formulary Committee. London, UK: BMJ Group and Pharmaceutical Press.
- (18) ATC/DDD Index 2014. 2014. World Health Organisation Collaborating Centre for Drug Statistics Methodology[Internet]. Oslo, Norway: World Health Organisation;[cited 2014 Nov 1]. Available from <http://www.whocc.no/>.
- (19) Khan NF, Perera R, Harper S, Rose PW. Adaptation and validation of the Charlson Index for Read/OXMIS coded databases. BMC Fam Pract 2010;11:1.
- (20) Noble M, Wright G, Dibben C, Smith GAN, McLennan D, Antilla C et al. Indices of deprivation 2004: Report to the office of the deputy prime minister. London, UK: Neighbourhood Renewal Unit; 2004.

- (21) Levesque LE, Hanley JA, Kezouh A, Suissa S. Problem of immortal time bias in cohort studies: example using statins for preventing progression of diabetes. *BMJ* 2010;340:b5087.
- (22) Chubak J, Boudreau DM, Wirtz HS, McKnight B, Weiss NS. Threats to Validity of Nonrandomized Studies of Postdiagnosis Exposures on Cancer Recurrence and Survival. *J Natl Cancer Inst* 2013;105:1456-1462.
- (23) Zhou Z, Rahme E, Abrahamowicz M, Pilote L. Survival bias associated with time-to-treatment initiation in drug effectiveness evaluation: a comparison of methods. *Am J Epidemiol* 2005;162:1016-1023.
- (24) Weinberg CR. Toward a clearer definition of confounding. *Am J Epidemiol* 1993;137:1-8.
- (25) Schisterman EF, Cole SR, Platt RW. Overadjustment bias and unnecessary adjustment in epidemiologic studies. *Epidemiology* 2009;20:488-495.
- (26) Han JY, Lee SH, Yoo NJ, Hyung LS, Moon YJ, Yun T et al. A randomized phase II study of gefitinib plus simvastatin versus gefitinib alone in previously treated patients with advanced non-small cell lung cancer. *Clin Cancer Res* 2011;17:1553-1560.
- (27) Phase I study of a statin + erlotinib for advanced solid malignancies with focus on squamous cell carcinomas and NSCLC. 2014. *ClinicalTrials.gov*[Internet]. Bethesda, USA: National Institutes of Health; [cited 2015 Nov 1]. Available from <http://clinicaltrials.gov/show/NCT00966472>.
- (28) BIBW 2992 plus simvastatin vs. BIBW 2992 in previously treated patients with advanced non-adenocarcinomatous NSCLC. *ClinicalTrials.gov*[Internet]. Bethesda, USA: National Institutes of Health; [cited 2015 Nov 1]. Available from <http://clinicaltrials.gov/show/NCT01156545>.
- (29) Irinotecan/cisplatin with or without simvastatin in chemo-naïve patients with extensive disease-small cell lung cancer. *ClinicalTrials.gov*[Internet]. Bethesda, USA: National

Institutes of Health; [cited 2015 Nov 1]. Available from

<http://clinicaltrials.gov/show/NCT01441349>.

(30) Irinotecan/cisplatin plus simvastatin in extensive disease-small cell lung cancer.

ClinicalTrials.gov[Internet]. Bethesda, USA: National Institutes of Health; [cited 2015 Nov

1]. Available from <http://clinicaltrials.gov/show/NCT00452634>.

(31) Stewart D, Cunningham IT, Hansford D, John D, McCaig D, McLay J. General practitioners' views and experiences of over-the-counter simvastatin in Scotland. *Br J Clin Pharmacol* 2010;70:356-359.

(32) Bosco JL, Silliman RA, Thwin SS, Geiger AM, Buist DS, Prout MN et al. A most stubborn bias: no adjustment method fully resolves confounding by indication in observational studies. *J Clin Epidemiol* 2010;63:64-74.

(33) Shrank WH, Patrick AR, Brookhart MA. Healthy user and related biases in observational studies of preventive interventions: a primer for physicians. *J Gen Intern Med* 2011;26:546-550.

Table 1. Characteristics of lung cancer patients by statin use before and after diagnosis.

Characteristics	Statin use in year prior to diagnosis <sup>a</sup>		Statin use in year after diagnosis <sup>b</sup>		Statin use after diagnosis <sup>b</sup>	
	Ever n (%)	Never (%)	User n (%)	Non-user (%)	User n (%)	Non-user (%)
	[n=13,398]		[n=3,638]		[n=3,638]	
Year of diagnosis: 1998-2000	102 (3)	2,204 (22)	24 (3)	618 (22)	79 (7)	563 (22)
2001-2003	451 (14)	2,893 (29)	132 (15)	754 (27)	217 (19)	669 (26)
2004-2006	1,118 (34)	2,650 (26)	279 (32)	690 (25)	347 (31)	622 (25)
2007-2009	1,616 (49)	2,362 (23)	428 (50)	713 (26)	472 (42)	669 (27)
Age at diagnosis: < 50	16 (0)	405 (4)	7 (1)	166 (6)	14 (1)	159 (6)
50-59	230 (7)	1,363 (13)	84 (10)	486 (18)	132 (12)	438 (17)
60-69	966 (29)	2,607 (26)	295 (34)	873 (31)	377 (34)	791 (31)
70-79	1,447 (44)	3,447 (34)	360 (42)	897 (32)	451 (40)	806 (32)
80-89	598 (18)	2,045 (20)	115 (13)	325 (12)	137 (12)	303 (12)
≥ 90	30 (1)	242 (2)	2 (0)	28 (1)	4 (0)	26 (1)
Gender: Males	2,030 (62)	5,886 (58)	541 (63)	1,563 (56)	690 (62)	1,414 (56)
Treatment within 6 months of cancer diagnosis						
Surgery <sup>c</sup>	339 (13)	883 (11)	212 (31)	659 (31)	333 (38)	538 (28)
Chemotherapy	743 (23)	2,287 (23)	267 (31)	993 (36)	324 (29)	936 (37)
Radiotherapy	1,028 (31)	3,197 (32)	319 (37)	1,062 (38)	377 (34)	1,004 (40)
Histology: Non-small cell	1,901 (58)	5,627 (56)	638 (74)	1,982 (71)	835 (75)	1,785 (71)
Small cell	418 (13)	1,257 (12)	89 (10)	335 (12)	108 (10)	316 (13)
Missing	968 (29)	3,225 (32)	136 (16)	458 (17)	172 (15)	422 (17)
Smoking status prior to cancer diagnosis						
Non-smoker	372 (11)	1,338 (13)	100 (12)	385 (14)	128 (11)	357 (14)
Ex-smoker	1,688 (51)	3,175 (31)	439 (51)	955 (34)	525 (47)	869 (34)
Current smoker	1,149 (35)	4,305 (43)	295 (34)	1,144 (41)	411 (37)	1,024 (41)
Missing	78 (2)	1,291 (13)	29 (3)	291 (10)	51 (5)	269 (11)
Alcohol consumption prior to diagnosis						
Never	564 (17)	1,271 (13)	132 (15)	340 (12)	165 (15)	307 (12)
Ever	2,175 (66)	5,709 (56)	586 (68)	1,671 (60)	750 (67)	1,507 (60)
Missing	548 (17)	3,129 (31)	145 (17)	764 (28)	200 (18)	709 (28)
BMI (kg/m <sup>2</sup> ) prior to diagnosis: mean (sd)	26.4 (5.0)	24.8 (4.8)	26.8 (4.9)	25.2 (4.7)	26.6 (4.9)	25.1 (4.7)
Underweight (<18.5)	119 (4)	510 (5)	23 (3)	109 (4)	29 (3)	103 (4)
Normal (18.5 to 25)	1,074 (33)	3,451 (34)	257 (30)	980 (35)	336 (30)	901 (36)
Overweight (25-30)	1,124 (34)	2,244 (22)	296 (34)	722 (26)	378 (34)	640 (25)
Obese (>30)	598 (18)	946 (9)	175 (20)	299 (11)	214 (19)	260 (10)
Missing	372 (11)	2,958 (27)	112 (13)	665 (24)	158 (14)	619 (25)
Deprivation fifth: 1 <sup>st</sup> (least deprived)	552 (17)	1,786 (17)	154 (18)	536 (19)	201 (18)	489 (19)
2 <sup>nd</sup>	638 (19)	1,945 (25)	160 (19)	527 (19)	204 (18)	483 (19)
3 <sup>rd</sup>	648 (20)	2,080 (21)	166 (19)	557 (20)	213 (19)	510 (20)
4 <sup>th</sup>	712 (22)	2,194 (22)	185 (21)	621 (22)	243 (22)	563 (22)
5 <sup>th</sup> (most deprived)	725 (22)	2,070 (20)	195 (23)	528 (19)	251 (23)	472 (19)
Missing	12 (0)	33 (0)	3 (0)	6 (0)	3 (0)	6 (0)
Comorbidity prior to cancer diagnosis						
Cerebrovascular disease	552 (17)	697 (7)	106 (12)	140 (5)	125 (11)	121 (5)
Chronic pulmonary disease	871 (27)	2,517 (25)	226 (26)	703 (25)	296 (27)	633 (25)
Congestive heart disease	307 (9)	593 (6)	64 (7)	98 (4)	72 (6)	90 (4)
Diabetes	854 (12)	600 (6)	203 (24)	132 (5)	230 (21)	105 (4)
Myocardial infarction	817 (25)	368 (4)	186 (22)	106 (4)	210 (19)	82 (3)
Peptic ulcer disease	263 (8)	743 (7)	58 (7)	188 (7)	79 (7)	167 (7)
Peripheral vascular disease	695 (21)	677 (7)	173 (20)	161 (6)	194 (17)	140 (6)
Renal disease	480 (15)	358 (4)	98 (11)	99 (4)	111 (10)	86 (3)
Other medication use after diagnosis						
Low dose aspirin use <sup>d</sup>	2,109 (64)	1,770 (18)	550 (64)	620 (22)	710 (64)	460 (18)
Beta-blocker use <sup>d</sup>	1,050 (32)	1,090 (11)	310 (36)	378 (14)	407 (37)	281 (11)

<sup>a</sup> Analysis includes lung cancer patients who have more than 1 year of records prior to diagnosis. <sup>b</sup> Analysis includes lung cancer patients who live more than 1 year after diagnosis. <sup>c</sup> Excluding cancer patients from Thames Registry as surgery information not available. <sup>d</sup> Low dose aspirin and beta-blocker use ever after diagnosis for statin use after diagnosis columns, low dose aspirin and beta-blocker use in year prior to diagnosis for statin use in year prior to diagnosis column.

Table 2. Association between statin usage after cancer diagnosis and cancer-specific and all-cause mortality in lung cancer patients.

Medication usage after diagnosis	Cancer-specific mortality	All-cause mortality	All patients	Person years	Cancer-specific mortality				All-cause mortality			
					Unadjusted HR (95%CI)	P	Adjusted <sup>a</sup> HR (95%CI)	P	Unadjusted HR (95%CI)	P	Adjusted <sup>a</sup> HR (95%CI)	P
<i>Number of patients</i>					<i>[n=3,638]</i>		<i>[n=2,792]</i>		<i>[n=3,638]</i>		<i>[n=2,792]</i>	
Statin non-user	1,669	1,962	2,523	4,619	1.00		1.00		1.00		1.00	
Statin user <sup>b</sup>	556	700	1,115	2,135	0.84 (0.76, 0.92)	<0.001	0.89 (0.78, 1.02)	0.09	0.89 (0.82, 0.97)	0.01	0.91 (0.80, 1.02)	0.10
Statin user 1 to 11 prescriptions <sup>c</sup>	355	422	535	977	0.84 (0.75, 0.94)	0.003	0.94 (0.81, 1.09)	0.39	0.88 (0.79, 0.98)	0.02	0.95 (0.83, 1.09)	0.44
Statin user ≥ 12 prescriptions <sup>c</sup>	201	278	580	1,158	0.84 (0.72, 0.97)	0.02	0.81 (0.67, 0.98)	0.03	0.91 (0.80, 1.04)	0.17	0.84 (0.71, 0.99)	0.04
Statin user 1 to 365 ddds <sup>c</sup>	337	406	493	903	0.90 (0.80, 1.01)	0.07	0.94 (0.81, 1.09)	0.40	0.95 (0.85, 1.06)	0.34	0.96 (0.84, 1.10)	0.57
Statin user ≥ 365 ddds <sup>c</sup>	219	294	622	1,232	0.76 (0.66, 0.88)	<0.001	0.82 (0.68, 0.98)	0.03	0.82 (0.72, 0.93)	0.002	0.83 (0.70, 0.97)	0.02
Simvastatin non-user	1,912	2,227	2,867	5,210	1.00		1.00		1.00		1.00	
Simvastatin user <sup>b</sup>	343	435	771	1,544	0.77 (0.68, 0.86)	<0.001	0.80 (0.69, 0.93)	0.003	0.80 (0.72, 0.89)	<0.001	0.81 (0.71, 0.92)	0.001
Simvastatin 1 to 11 prescriptions <sup>c</sup>	220	268	387	777	0.75 (0.65, 0.86)	<0.001	0.82 (0.69, 0.98)	0.03	0.78 (0.69, 0.89)	<0.001	0.84 (0.72, 0.98)	0.03
Simvastatin ≥ 12 prescriptions <sup>c</sup>	123	167	384	767	0.80 (0.67, 0.97)	0.02	0.76 (0.60, 0.95)	0.01	0.84 (0.72, 0.99)	0.04	0.75 (0.62, 0.91)	0.004
Atorvastatin non-user	2,083	2,432	3,301	6,126	1.00		1.00		1.00		1.00	
Atorvastatin user <sup>b</sup>	172	230	337	628	0.93 (0.80, 1.09)	0.37	0.93 (0.78, 1.12)	0.47	1.04 (0.91, 1.20)	0.52	1.03 (0.88, 1.21)	0.68
Pravastatin non-user	2,216	2,615	3,564	6,620	1.00		1.00		1.00		1.00	
Pravastatin user <sup>b</sup>	39	47	74	133	1.02 (0.74, 1.39)	0.93	1.23 (0.85, 1.77)	0.27	1.01 (0.76, 1.35)	0.94	1.11 (0.79, 1.56)	0.53
Rosuvastatin non-user	2,237	2,640	3,598	6,668	1.00		1.00		1.00		1.00	
Rosuvastatin user <sup>b</sup>	18	22	40	86	0.93 (0.58, 1.47)	0.74	1.15 (0.66, 2.01)	0.61	0.89 (0.58, 1.35)	0.58	0.87 (0.51, 1.48)	0.61
Fluvastatin non-user	2,246	2,651	3,624	6,726	1.00		1.00		1.00		1.00	
Fluvastatin user <sup>b</sup>	9	11	14	28	1.16 (0.60, 2.23)	0.66	0.98 (0.47, 2.09)	0.98	1.17 (0.65, 2.12)	0.60	1.07 (0.55, 2.06)	0.85
Any lipophilic <sup>d</sup> non-user	1,901	2,214	2,851	1,573	1.00		1.00		1.00		1.00	
Any lipophilic statin user <sup>b</sup>	354	448	787	5,181	0.78 (0.69, 0.87)	<0.001	0.81 (0.70, 0.94)	0.01	0.81 (0.73, 0.90)	<0.001	0.82 (0.72, 0.93)	0.002
Any hydrophilic <sup>d</sup> non-user	2,027	2,367	3,205	5,945	1.00		1.00		1.00		1.00	
Any hydrophilic statin user <sup>b</sup>	228	295	433	809	0.96 (0.84, 1.10)	0.54	1.00 (0.85, 1.18)	0.97	1.04 (0.92, 1.17)	0.54	1.05 (0.91, 1.22)	0.49

<sup>a</sup> Adjusted for year of diagnosis, age at diagnosis, gender, radiotherapy within 6 months, chemotherapy within 6 months, surgery within 6 months, comorbidities (prior to diagnosis, including cerebrovascular disease, chronic pulmonary disease, congestive heart disease, diabetes, myocardial infarction, peptic ulcer disease, peripheral vascular disease, renal disease), other medication use (after diagnosis, as time varying covariates, specifically low dose aspirin and beta-blockers) and deprivation (in fifths).

<sup>b</sup> Medication use modelled as a time varying covariate with an individual considered a non-user prior to 6 months after first medication usage and a user after this time, excludes deaths in the year after cancer diagnosis.

<sup>c</sup> Medication use modelled as a time varying covariate with an individual considered a non-user prior to 6 months after first medication usage, a user of 0 to 12 prescriptions from 6 months after first prescription to 6 months after 12<sup>th</sup> prescription (or 365<sup>th</sup> DDD) and a greater user after this time, excludes deaths in the year after cancer diagnosis.

<sup>d</sup> Lipophilic statins include simvastatin, fluvastatin, cerivastatin; hydrophilic statins include atorvastatin, pravastatin and rosuvastatin.

Table 3. Sensitivity analyses for association between statin use and cancer-specific mortality in lung cancer patients.

	Cancer-specific mortality	All patients	Person years	User versus non-user		1-11 prescriptions v. none		≥ 12 prescriptions v. none	
				Adj HR <sup>a</sup> (95% CI)	P	Adj HR <sup>a</sup> (95% CI)	P	Adj HR <sup>a</sup> (95% CI)	P
Main analysis: Statin user versus non-user after diagnosis	1,735	2,792	5,227	0.89 (0.78, 1.02)	0.09	0.94 (0.81, 1.09)	0.39	0.81 (0.67, 0.98)	0.03
Sub group analyses: Statin user versus non-user after diagnosis, restricted to:									
Males	1,007	1,620	2,928	0.94 (0.79, 1.12)	0.62	0.99 (0.82, 1.21)	0.96	0.85 (0.66, 1.09)	0.19
Females	728	1,172	2,299	0.83 (0.67, 1.03)	0.09	0.86 (0.68, 1.10)	0.23	0.78 (0.58, 1.06)	0.11
Pre-diagnosis statin non-users <sup>b</sup>	1,222	1,912	3,691	0.78 (0.60, 1.02)	0.07	0.64 (0.45, 0.91)	0.01	1.06 (0.72, 1.57)	0.77
Pre-diagnosis statin users <sup>b</sup>	402	697	1,132	0.76 (0.54, 1.07)	0.12	0.83 (0.59, 1.17)	0.29	0.63 (0.42, 0.93)	0.02
Small cell lung cancer	268	338	489	0.66 (0.44, 0.99)	0.05	0.64 (0.41, 0.99)	0.05	0.74 (0.41, 1.36)	0.33
Non-small cell lung cancer	1,163	1,995	4,090	0.93 (0.79, 1.09)	0.38	1.01 (0.84, 1.21)	0.90	0.80 (0.64, 1.01)	0.06
Sensitivity analyses: Statin user versus non-user after diagnosis									
Increasing lag to 1 year	1,735	2,792	5,227	0.92 (0.80, 1.05)	0.23	0.92 (0.80, 1.07)	0.30	0.90 (0.72, 1.14)	0.39
Increasing lag to 1 years (simvastatin only)	1,735	2,792	5,227	0.86 (0.74, 1.00)	0.06	0.84 (0.71, 1.00)	0.05	0.91 (0.70, 1.19)	0.50
Additionally adjusting for smoking prior to diagnosis	1,584	2,564	4,760	0.90 (0.78, 1.03)	0.12	0.95 (0.82, 1.11)	0.52	0.82 (0.66, 0.97)	0.03
Additionally adjusting for BMI prior to diagnosis	1,340	2,206	4,154	0.93 (0.80, 1.08)	0.34	0.98 (0.83, 1.15)	0.80	0.85 (0.68, 1.04)	0.12
Additionally adjusting for small cell/non-small cell	1,431	2,333	4,579	0.86 (0.75, 1.00)	0.05	0.92 (0.78, 1.09)	0.32	0.77 (0.62, 0.95)	0.02
Additionally adjusting for bisphosphonate use <sup>c</sup>	1,735	2,792	5,227	0.89 (0.78, 1.02)	0.09	0.94 (0.81, 1.09)	0.40	0.81 (0.67, 0.98)	0.03
Based upon first year after diagnosis <sup>d</sup>	1,735	2,792	5,227	0.90 (0.78, 1.03)	0.13	0.97 (0.83, 1.12)	0.66	0.75 (0.60, 0.93)	0.01
Nested case-control analysis <sup>e</sup>	1,705			0.92 (0.79, 1.07)	0.27	0.98 (0.82, 1.15)	0.77	0.81 (0.65, 1.01)	0.06

<sup>a</sup> Except where otherwise stated, all analyses of post-diagnostic statin use adjusted for year of diagnosis, age at diagnosis, gender, surgery within 6 months of diagnosis, radiotherapy within 6 months, chemotherapy within 6 months, comorbidities (prior to diagnosis, including cerebrovascular disease, chronic pulmonary disease, congestive heart disease, diabetes, myocardial infarction, peptic ulcer disease, peripheral vascular disease, renal disease), other medication use (after diagnosis, as time varying covariates, specifically low dose aspirin and beta-blockers) and deprivation (in fifths).

<sup>b</sup> Based upon statin use in the year prior to diagnosis, restricted to individuals with 1 year of records prior to lung cancer diagnosis.

<sup>c</sup> Contains all confounders in <sup>a</sup> along with bisphosphonate use after diagnosis (as a time varying covariate).

<sup>d</sup> Simplified analysis, not requiring time varying covariate use, comparing statin users to statin non-users in the first year after diagnosis in individuals living more than 1 year after cancer diagnosis, adjusted for all confounders in <sup>a</sup> but other medication use also restricted to first year after diagnosis.

<sup>e</sup> Estimate and 95% CIs are for adjusted OR, 25% (433/1705) of cancer-specific deaths used statins compared with 29% (2388/8150) of risk-set controls (not dying from cancer), matched on age at diagnosis, year of diagnosis, gender and adjusted for all other confounders in <sup>a</sup>.



Table 4. Association between statin usage in the year prior to diagnosis and cancer-specific death in lung cancer patients.

Medication usage in year prior to diagnosis	Cancer-specific mortality	All-cause mortality	All patients	Person years	Cancer-specific mortality				All-cause mortality			
					Unadjusted HR (95%CI)	P	Adjusted <sup>a</sup> HR (95%CI)	P	Unadjusted HR (95%CI)	P	Adjusted <sup>a</sup> HR (95%CI)	P
<i>Number of patients</i>					<i>[n=13,398]</i>		<i>[n=13,353]</i>		<i>[n=13,398]</i>		<i>[n=13,353]</i>	
Statin non-user	8,064	8,926	10,111	9430	1.00		1.00		1.00		1.00	
Statin user	2,558	2,898	3,287	3,077	0.94 (0.90, 0.98)	0.007	0.88 (0.83, 0.93)	<0.001	0.97 (0.93, 1.01)	0.10	0.89 (0.85, 0.94)	<0.001
Statin user 1 to 11 prescriptions	1780	2,011	2,290	2,235	0.92 (0.87, 0.96)	0.001	0.87 (0.82, 0.92)	<0.001	0.94 (0.89, 0.98)	0.01	0.88 (0.83, 0.93)	<0.001
Statin user ≥ 12 prescriptions	778	887	997	841	1.00 (0.93, 1.08)	0.93	0.91 (0.84, 0.99)	0.02	1.04 (0.97, 1.11)	0.29	0.92 (0.85, 1.00)	0.04
Statin user 1 to 365 ddds	1,737	1,981	2,235	2,110	0.94 (0.89, 0.99)	0.02	0.88 (0.83, 0.93)	<0.001	0.97 (0.93, 1.02)	0.27	0.89 (0.84, 0.95)	<0.001
Statin user ≥ 365 ddds	821	917	1,052	967	0.94 (0.87, 1.01)	0.08	0.89 (0.82, 0.96)	0.004	0.95 (0.89, 1.02)	0.14	0.88 (0.82, 0.95)	0.002
Simvastatin non-user	8,977	9,966	11,265	10,499	1.00		1.00		1.00		1.00	
Simvastatin user	1,645	1,858	2,133	2,008	0.95 (0.90, 1.00)	0.05	0.92 (0.86, 0.97)	0.004	0.97 (0.92, 1.02)	0.18	0.92 (0.87, 0.97)	0.003
Atorvastatin non-user	9,832	10,922	12,392	11,549	1.00		1.00		1.00		1.00	
Atorvastatin user	790	902	1,006	958	0.93 (0.87, 1.00)	0.06	0.91 (0.85, 0.99)	0.02	0.96 (0.90, 1.03)	0.28	0.93 (0.87, 1.00)	0.05
Pravastatin non-user	10,468	11,643	13,200	12,332	1.00		1.00		1.00		1.00	
Pravastatin user	154	181	198	174	0.99 (0.84, 1.16)	0.88	0.96 (0.82, 1.12)	0.60	1.05 (0.90, 1.21)	0.54	1.00 (0.86, 1.16)	0.96
Rosuvastatin non-user	10,555	11,750	13,310	12,418	1.00		1.00		1.00		1.00	
Rosuvastatin user	67	74	88	89	0.83 (0.66, 1.06)	0.14	0.82 (0.65, 1.05)	0.11	0.83 (0.66, 1.05)	0.12	0.81 (0.64, 1.02)	0.07
Fluvastatin non-user	10,575	11,771	13,341	12,464	1.00		1.00		1.00		1.00	
Fluvastatin user	47	53	57	43	1.03 (0.78, 1.38)	0.82	1.01 (0.76, 1.35)	0.93	1.06 (0.81, 1.39)	0.68	1.03 (0.78, 1.35)	0.85
Any lipophilic <sup>b</sup> non-user	8,922	9,904	11,200	10,455	1.00		1.00		1.00		1.00	
Any lipophilic statin user	1,700	1,920	2,198	2,052	0.95 (0.91, 1.00)	0.07	0.92 (0.87, 0.98)	0.01	0.97 (0.93, 1.02)	0.25	0.92 (0.87, 0.98)	0.01
Any hydrophilic <sup>b</sup> non-user	9,631	10,690	12,129	11,302	1.00		1.00		1.00		1.00	
Any hydrophilic statin user	991	1,134	1,269	1,205	0.93 (0.87, 0.99)	0.02	0.90 (0.84, 0.96)	0.003	0.96 (0.90, 1.02)	0.17	0.92 (0.86, 0.98)	0.01

<sup>a</sup> Adjusted for year of diagnosis, age at diagnosis, gender, comorbidities (prior to diagnosis, including cerebrovascular disease, chronic pulmonary disease, congestive heart disease, diabetes, myocardial infarction, peptic ulcer disease, peripheral vascular disease, renal disease), other medication use (in year prior to diagnosis, specifically low dose aspirin and beta-blockers) and deprivation (in fifths).

<sup>b</sup> Lipophilic statins include simvastatin, fluvastatin, cerivastatin; hydrophilic statins include atorvastatin, pravastatin and rosuvastatin.

Table 5. Sensitivity analyses for association between statin use and cancer-specific mortality in lung cancer patients.

	Cancer-specific mortality	All patients	Person years	Unadjusted HR (95% CI)	P	Adjusted <sup>a</sup> HR (95% CI)	P
Main analysis: Pre-diagnostic statin use <sup>b</sup>	10,622	13,398	12,507	0.94 (0.90, 0.98)	0.007	0.88 (0.83, 0.93)	<0.001
Subgroup analyses							
Male	6,292	7,918	7,115	0.91 (0.86, 0.96)	0.001	0.87 (0.80, 0.93)	<0.001
Female	4,330	5,480	5,391	0.99 (0.92, 1.06)	0.80	0.91 (0.83, 0.99)	0.02
Sensitivity analyses							
Smoking prior to diagnosis available (and adjusted for)	9,519	9,519	11,329	0.96 (0.92, 1.00)	0.07	0.89 (0.84, 0.95)	<0.001
BMI prior to diagnosis available (and adjusted for)	7,959	10,040	9,783	0.98 (0.93, 1.03)	0.42	0.92 (0.87, 0.98)	0.01
Additionally adjusting for bisphosphonate use <sup>c</sup>	10,622	13,398	12,507	0.94 (0.90, 0.98)	0.007	0.88 (0.83, 0.93)	<0.001
Statin use between 2 years and 6 months prior to diagnosis <sup>d</sup>	9,733	12,260	11,437	0.93 (0.89, 0.98)	0.01	0.87 (0.82, 0.92)	<0.001

<sup>a</sup> Except where otherwise stated, adjusted for year of diagnosis, age at diagnosis, gender, comorbidities (prior to diagnosis, including cerebrovascular disease, chronic pulmonary disease, congestive heart disease, diabetes, myocardial infarction, peptic ulcer disease, peripheral vascular disease, renal disease), other medication use (in year prior to diagnosis, specifically low dose aspirin and beta-blockers) and deprivation (in fifths).

<sup>b</sup> Based upon use in the year prior to diagnosis, restricted to individuals with 1 year of records prior to lung cancer diagnosis.

<sup>c</sup> Contains all confounders in <sup>a</sup> along with bisphosphonate use in year prior to diagnosis.

<sup>d</sup> Restricted to individuals with 2 years of records prior to diagnosis, removing prescriptions in the 6 months prior to lung cancer diagnosis as these could reflect increased medical care due to early symptoms.